

# **Pelvic Pain Treatment with PEMF Magnetic Therapy**

## **Electrochemical therapy of pelvic pain: effects of pulsed electromagnetic fields (PEMF) on Pelvic Pain & tissue trauma.**

Jorgensen WA,  
Frome BM,  
Wallach C.

International Pain Research Institute, Los Angeles, California.

Unusually effective and long-lasting relief of pelvic pain of gynaecological origin has been obtained consistently by short exposures of affected areas to the application of a magnetic induction device producing short, sharp, magnetic-field pulses of a minimal amplitude to initiate the electrochemical phenomenon of electroporation within a 25 cm<sup>2</sup> focal area. Treatments are short, fast-acting, economical and in many instances have obviated surgery. This report describes typical cases such as dysmenorrhoea, endometriosis, ruptured ovarian cyst, acute lower urinary tract infection, post-operative haematoma, and persistent dyspareunia in which pulsed magnetic field treatment has not, in most cases, been supplemented by analgesic medication. Of 17 female patients presenting with a total of 20 episodes of pelvic pain, of which 11 episodes were acute, seven chronic and two acute as well as chronic, 16 patients representing 18 episodes (90%) experienced marked, even dramatic relief, while two patients representing two episodes reported less than complete pain relief.

J Rheumatol. 1993 Mar;20(3):456-60.